

GENERALIZATION OF THE HAUSDORFF MOMENT PROBLEM

DAVID BORWEIN AND AMNON JAKIMOVSKI

1. Introduction. Suppose throughout that $\{k_n\}$ is a sequence of positive integers, that

$$0 \leq l_0 < l_1 < l_2 < \dots < l_n, l_n \rightarrow \infty, \sum_{n=1}^{\infty} \frac{k_n}{l_n} = \infty,$$

that $k_0 = 1$ if $l_0 = 1$, and that $\{u_n^{(r)}\}$ ($r = 0, 1, \dots, k_n - 1, n = 0, 1, \dots$) is a sequence of real numbers. We shall be concerned with the problem of establishing necessary and sufficient conditions for there to be a function α satisfying

$$(1) \quad (-1)^r u_n^{(r)} = \int_0^1 t^{l_n} \log^r t \, d\alpha(t)$$

for $r = 0, 1, \dots, k_n - 1, n = 0, 1, \dots$

and certain additional conditions. The case $l_0 = 0, k_n = 1$ for $n = 0, 1, \dots$ of the problem is the version of the classical moment problem considered originally by Hausdorff [5], [6], [7]; the above formulation will emerge as a natural generalization thereof. An alternative formulation of the problem is to express it as the "infinite Hermite interpolation problem" of establishing necessary and sufficient conditions for a function F to be a Laplace transform of the form

$$F(z) = \int_0^{\infty} e^{-uz} d\gamma(u)$$

and to satisfy

$$F^{(r)}(l_n) = (-1)^r u_n^{(r)} \text{ for } r = 0, 1, \dots, k_n - 1, n = 0, 1, \dots$$

Considerable simplification is obtained by adoption of the following notation. Construct a monotonic sequence $\{\lambda_s\}$ from $\{l_n\}$ by repeating each l_n k_n times. Then

$$0 \leq \lambda_0 \leq \lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_n, \lambda_1 > 0, \lambda_n \rightarrow \infty, \sum_{n=1}^{\infty} \frac{1}{\lambda_n} = \infty.$$

Received March 5, 1980 and in revised form January 5, 1981. This research was supported in part by the Natural Sciences and Engineering Research Council of Canada Grant A-2983.

For each s there is an integer $n(s)$ such that $\lambda_s = l_{n(s)}$. Let $m_s = k_{n(s)}$ and construct a sequence $\{\mu_s^{(r)}\}$ ($r = 0, 1, \dots, m_s - 1, s = 0, 1, \dots$) from $\{u_n^{(r)}\}$ by setting $\mu_s^{(r)} = u_{n(s)}^{(r)}$. Then m_s is the multiplicity of λ_s , i.e., it is the number of indices j for which $\lambda_j = \lambda_s$; and $\mu_j^{(r)} = \mu_s^{(r)}$ whenever $\lambda_j = \lambda_s$. Formula (1) can be written in the equivalent form

$$(2) \quad (-1)^r \mu_s^{(r)} = \int_0^1 t^{\lambda_s} \log^r t \, d\alpha(t)$$

for $r = 0, 1, \dots, m_s - 1, s = 0, 1, \dots$

For $0 \leq k \leq s \leq n$, let $m_s(k, n)$ be the multiplicity of λ_s among $\lambda_k, \lambda_{k+1}, \dots, \lambda_n$. By a standard result on Hermite interpolation (see [3, p. 29]) there is a unique polynomial $P_n(z)$ of degree at most n such that

$$(3) \quad P_n^{(r)}(\lambda_s) = (-1)^r \mu_s^{(r)} \text{ for } r = 0, 1, \dots, m_s(0, n) - 1, \\ s = 0, 1, \dots, n.$$

It is known (see [11, p. 45]) that

$$P_n(z) = \sum_{k=0}^n u[\lambda_k, \dots, \lambda_n] (\lambda_{k+1} - z) \dots (\lambda_n - z)$$

where the divided difference $u[\lambda_k, \dots, \lambda_n]$ is given by

$$u[\lambda_k, \dots, \lambda_n] = -\frac{1}{2\pi i} \int_{C_{kn}} \frac{P_n(z) dz}{c_{kn} (\lambda_k - z) \dots (\lambda_n - z)},$$

C_{kn} being a positively sensed Jordan contour enclosing $\lambda_k, \lambda_{k+1}, \dots, \lambda_n$. For $0 \leq k \leq n, 0 < t \leq 1$, let

$$\lambda_{nk} = \lambda_{k+1} \dots \lambda_n u[\lambda_k, \dots, \lambda_n],$$

$$(4) \quad \lambda_{nk}(t) = -\lambda_{k+1} \dots \lambda_n \frac{1}{2\pi i} \int_{C_{kn}} \frac{t^z dz}{(\lambda_k - z) \dots (\lambda_n - z)},$$

$$\lambda_{nk}(0) = \lambda_{nk}(0+),$$

with the convention that products such as $\lambda_{k+1} \dots \lambda_n = 1$ when $k = n$.

If $f(z)$ is analytic inside and on C_{kn} then, by the theory of residues,

$$\int_{C_{kn}} \frac{f(z) dz}{(\lambda_k - z) \dots (\lambda_n - z)}$$

is a linear combination, with coefficients depending only on $\lambda_k, \lambda_{k+1}, \dots, \lambda_n$, of the values $f^{(r)}(\lambda_s), r = 0, 1, \dots, m_s(k, n) - 1, s = k, k + 1, \dots, n$. It follows that $\lambda_{nk}(t)$ is a linear combination of the functions $t^{\lambda_s} \log^r t, r = 0, 1, \dots, m_s(k, n) - 1, s = k, k + 1, \dots, n$ and that λ_{nk} is the same linear combination with $(-1)^r \mu_s^{(r)}$ substituted for $t^{\lambda_s} \log^r t$. Consequently, if $\alpha \in BV$, where BV is the space of norma-

lized functions of bounded variation on $[0, 1]$, i.e., $\alpha(0) = 0$, $2\alpha(t) = \alpha(t+) + \alpha(t-)$ for $0 < t < 1$, and if

$$(-1)^r \mu_s^{(r)} = \int_0^1 t^{\lambda_s} \log^r t \, d\alpha(t) \quad \text{for } 0 \leq r < m_s(k, n), \quad k \leq s \leq n,$$

then

$$\lambda_{nk} = \int_0^1 \lambda_{nk}(t) d\alpha(t).$$

An explicit formula for $u[\lambda_k, \dots, \lambda_n]$ can be obtained by evaluating

$$\frac{1}{2\pi i} \int_{c_{kn}} \frac{t^2 dz}{(\lambda_k - z) \dots (\lambda_n - z)}$$

and substituting $(-1)^r \mu_s^{(r)}$ for $t^{\lambda_s} \log^r t$ in the result.

Let

$$D_0 = (1 + \lambda_0)d_0 = 1, \quad D_n = \left(1 + \frac{1}{\lambda_1}\right) \dots \left(1 + \frac{1}{\lambda_n}\right) \\ = (1 + \lambda_n)d_n \quad \text{for } n \geq 1.$$

Then, for $n \geq 0$,

$$D_n = \lambda_{n+1}d_{n+1} = \frac{\lambda_0}{1 + \lambda_0} + \sum_{k=0}^n d_k,$$

and, for $n > k \geq 0$,

$$(5) \quad \sum_{j=k+1}^n \frac{1}{1 + \lambda_j} = \sum_{j=k+1}^n \frac{d_j}{D_j} \leq \sum_{j=k+1}^n \int_{D_{j-1}}^{D_j} \frac{dx}{x} = \log \frac{D_n}{D_k} \\ \leq \sum_{j=k+1}^n \frac{d_j}{D_{j-1}} = \sum_{j=k+1}^n \frac{1}{\lambda_j}.$$

Further, it is known that if all the λ_n 's are different, then

$$(6) \quad 0 \leq \lambda_{ns}(t) \leq \sum_{k=0}^n \lambda_{nk}(t) \leq 1 \quad \text{for } 0 \leq t \leq 1, \quad 0 \leq s \leq n,$$

by [10, Lemma 1] and

$$(7) \quad \int_0^1 \lambda_{nk}(t) dt = \frac{d_k}{D_n} \quad \text{for } 0 \leq k \leq n$$

by [6, p. 294]. A simple continuity argument applied to (4) shows that (6) and (7) remain valid when different λ_n 's are allowed to coalesce.

Let θ be an even continuous convex function such that $\theta(u)/u \rightarrow 0$ as $u \rightarrow 0$ and $\theta(u)/u \rightarrow \infty$ as $u \rightarrow \infty$. Associated with this function is the Orlicz class L_θ of all functions x Lebesgue integrable over $[0, 1]$ for which

$$\int_0^1 \theta(x(t)) dt < \infty.$$

Let L_∞ be the space of measurable functions x on $[0, 1]$ with finite norm

$$\|x\|_\infty = \text{ess. sup}_{0 < t < 1} |x(t)|.$$

Let

$$M_\theta(n) = \sum_{k=0}^n \frac{d_k}{D_n} \theta\left(\frac{D_n}{d_k} \lambda_{nk}\right),$$

$$M_1(n) = \sum_{k=0}^n |\lambda_{nk}|,$$

$$M_\infty(n) = \max_{0 \leq k \leq n} |\lambda_{nk}| \frac{D_n}{d_k},$$

and let

$$M_\theta = \sup_{n \geq 0} M_\theta(n), \quad M_1 = \sup_{n \geq 0} M_1(n), \quad M_\infty = \sup_{n \geq 0} M_\infty(n).$$

The following two theorems are the main results established in the present paper.

THEOREM 1. *A necessary and sufficient condition for there to be a function*

- (i) $\alpha \in \text{BV}$ satisfying (1) is that $M_1 < \infty$;
- (ii) $\beta \in L_\infty$ satisfying

$$(8) \quad (-1)^r u_n^{(r)} = \int_0^1 t^{\lambda_n} \log^r t \beta(t) dt$$

for $r = 0, 1, \dots, k_n - 1, n = 0, 1, \dots$

is that $M_\infty < \infty$;

- (iii) $\beta \in L_\theta$ satisfying (8) is that $M_\theta < \infty$.

Furthermore

(iv) if (1) is satisfied by a function $\alpha \in \text{BV}$, then $M_1 = \int_0^1 |\alpha(t)| - \delta |\alpha(0+)|$ where $\delta = 0$ when $l_0 = 0$, $\delta = 1$ when $l_0 > 0$; moreover α is unique when $l_0 = 0$, and when $l_0 > 0$ it differs by a constant, over the interval $0 < t \leq 1$, from any other function in BV satisfying (1);

(v) if (8) is satisfied by a function $\beta \in L_\infty$, then β is essentially unique and $M_\infty = \|\beta\|_\infty$;

(vi) if (8) is satisfied by a function $\beta \in L_\theta$, then β is essentially unique and

$$M_\theta = \int_0^1 \theta(\beta(t)) dt.$$

THEOREM 2. For $n = 0, 1, \dots$,

$$M_1(n) \leq M_1(n+1), \quad M_\infty(n) \leq M_\infty(n+1), \quad M_\theta(n) \leq M_\theta(n+1);$$

and

$$\lim_{n \rightarrow \infty} M_1(n) = M_1, \lim_{n \rightarrow \infty} M_\infty(n) = M_\infty, \lim_{n \rightarrow \infty} M_\theta(n) = M_\theta.$$

The case $l_0 = 0$, $k_n = 1$ for $n = 0, 1, \dots$ of Theorem 1(i) was established by Hausdorff [5], [6] and Schoenberg [13] subsequently gave a different proof; the case $l_0 > 0$, $k_n = 1$ for $n = 0, 1, \dots$ was proved by Leviatan [9]. (See also [4].)

The case $l_n = n$, $k_n = 1$ for $n = 0, 1, \dots$ of Theorem 1(ii) is due to Hausdorff [7].

The case $l_n = n$, $k_n = 1$ for $n = 0, 1, \dots$, $\theta(u) = |u|^p$, $1 < p < \infty$, of Theorem 1(iii) is due to Hausdorff [7] and the case $k_n = 1$ for $n = 0, 1, \dots$ to Leviatan [9], [10]. (See also [1] and [2].)

See [2] and the references there given for known special cases of Theorem 2.

2. Preliminary results.

LEMMA 1. Let r, a be non-negative integers, let $0 < \lambda < \lambda_{a+1}$, and let

$$\delta_{nk} = \left(1 - \frac{\lambda}{\lambda_{k+1}}\right) \dots \left(1 - \frac{\lambda}{\lambda_n}\right) \left(\sum_{j=k+1}^n \frac{1}{\lambda_j - \lambda}\right)^r.$$

Then (i) δ_{nk} is uniformly bounded for $n > k \geq a$,

(ii) $\lim_{n \rightarrow \infty} \delta_{nk} = 0$ for $k \geq a$,

(iii) $\delta_{nk} - \left(\frac{D_k}{D_n}\right)^\lambda \log^r \frac{D_n}{D_k} \rightarrow 0$ uniformly when $n > k \rightarrow \infty$.

Proof. Let $0 < \epsilon < \lambda$, $\alpha = \lambda - \epsilon$, $\beta = \lambda + \epsilon$, let

$$\gamma = \gamma_{nk} = \sum_{j=k+1}^n \frac{1}{\lambda_j},$$

and, for $n > a$, let

$$u_n = 1 - \frac{\lambda}{\lambda_n} = e^{-\alpha n / \lambda_n}, v_n = \left(1 + \frac{1}{\lambda_n}\right)^{-\lambda} = e^{-\beta n / \lambda_n}.$$

Then $\alpha_n \rightarrow \lambda$, $\beta_n \rightarrow \lambda$ and so we can choose a positive integer $N \geq a$ so large that

$$|\alpha_n - \lambda| < \epsilon, |\beta_n - \lambda| < \epsilon \quad \text{for } n > N.$$

First, for $n > k \geq N$, we have that

$$0 < \delta_{nk} = u_{k+1} \dots u_n \left(\sum_{j=k+1}^n \frac{1}{\lambda_j - \lambda}\right)^r \leq e^{-\alpha \gamma} \gamma^r \left(\frac{\lambda_{k+1}}{\lambda_{k+1} - \lambda}\right)^r.$$

Since $\gamma_{nk} \rightarrow \infty$ as $n \rightarrow \infty$, it follows that (i) and (ii) hold for $k \geq N$. The extension of these conclusions to the range $N > k \geq a$ is simple.

Next, let

$$a_{nk} = |u_{k+1} \dots u_n - v_{k+1} \dots v_n| \left(\sum_{j=k+1}^n \frac{1}{\lambda_j - \lambda}\right)^r,$$

$$b_{nk} = v_{k+1} \dots v_n \left\{ \left(\sum_{j=k+1}^n \frac{1}{\lambda_j - \lambda}\right)^r - \log^r \frac{D_n}{D_k} \right\}.$$

Then, for $n > k \geq N$, we have that

$$(9) \quad 0 \leq a_{nk} \leq (e^{-\alpha \gamma} - e^{-\beta \gamma}) \gamma^r \left(\frac{\lambda_{k+1}}{\lambda_{k+1} - \lambda}\right)^r$$

$$\leq \gamma(\beta - \alpha) e^{-\alpha \gamma} \gamma^r \left(\frac{\lambda_{k+1}}{\lambda_{k+1} - \lambda}\right)^r$$

$$\leq \frac{2\gamma \epsilon \gamma^r (r+1)!}{(\alpha \gamma)^{r+1}} \left(\frac{\lambda_{k+1}}{\lambda_{k+1} - \lambda}\right)^r = \frac{2(r+1)!}{(\lambda - \epsilon)^{r+1}} \left(\frac{\lambda_{k+1}}{\lambda_{k+1} - \lambda}\right)^r \epsilon,$$

and, by (5), that

$$(10) \quad 0 \leq b_{nk} \leq v_{k+1} \dots v_n \left(\sum_{j=k+1}^n \frac{1}{\lambda_j - \lambda}\right)^{r-1} \left(\sum_{j=k+1}^n \frac{1}{\lambda_j - \lambda} - \log \frac{D_n}{D_k}\right)$$

$$\leq v_{k+1} \dots v_n \left(\sum_{j=k+1}^n \frac{1}{\lambda_j - \lambda}\right)^{r-1} \left(\sum_{j=k+1}^n \frac{1}{\lambda_j - \lambda} - \frac{1}{1 + \lambda_j}\right)$$

$$\leq v_{k+1} \dots v_n \left(\sum_{j=k+1}^n \frac{1}{\lambda_j - \lambda}\right)^r \frac{\lambda + 1}{\lambda_{k+1}}$$

$$\leq e^{-\alpha \gamma} r \gamma^r \left(\frac{\lambda_{k+1}}{\lambda_{k+1} - \lambda}\right)^r \frac{\lambda + 1}{\lambda_{k+1}} \leq \frac{r r!}{(\lambda - \epsilon)^r} \left(\frac{\lambda_{k+1}}{\lambda_{k+1} - \lambda}\right)^r \frac{\lambda + 1}{\lambda_{k+1}}.$$

It follows from (9) that $a_{nk} \rightarrow 0$ uniformly when $n > k \rightarrow \infty$, and from (10) that $b_{nk} \rightarrow 0$ uniformly when $n > k \rightarrow \infty$. Since

$$\left| \delta_{nk} - \left(\frac{D_k}{D_n}\right)^\lambda \log^r \frac{D_n}{D_k} \right| \leq a_{nk} + b_{nk} \quad \text{for } n > k \geq N,$$

conclusion (iii) follows.

LEMMA 2. Let $\psi(t) = (\lambda_{k+1} - t) \dots (\lambda_n - t)$ where $0 \leq k < n$ and $0 < t < \lambda_{k+1}$, and let r be a positive integer. Then

$$\left| \psi^{(r)}(t) - (-1)^r \psi(t) \left(\sum_{j=k+1}^n \frac{1}{\lambda_j - t}\right)^r \right| \leq \frac{M \psi(t)}{\lambda_{k+1} - t} \left(\sum_{j=k+1}^n \frac{1}{\lambda_j - t}\right)^{r-1}$$

where M is a number independent of t, k and n .

Proof. The result is evidently true with $M = 0$ when $r = 1$. Suppose therefore that $r \geq 2$ and let

$$\gamma_j = \frac{1}{\lambda_j - t}.$$

As easy inductive argument shows that

$$\frac{\psi^{(r)}(t)}{\psi(t)} - (-1)^r \left(\sum_{j=k+1}^n \gamma_j \right)^r$$

is equal to a linear combination with constant coefficients of terms of the form

$$\left(\sum_{j=k+1}^n \gamma_j^{a_1} \right)^{b_1} \left(\sum_{j=k+1}^n \gamma_j^{a_2} \right)^{b_2} \dots \left(\sum_{j=k+1}^n \gamma_j^{a_m} \right)^{b_m}$$

where the a_i 's and b_i 's are positive integers, $a_1 > 1$ and

$$a_1 b_1 + a_2 b_2 + \dots + a_m b_m = r.$$

Each of the terms is no greater than

$$\begin{aligned} & \gamma_{k+1} \left(\sum_{j=k+1}^n \gamma_j^{a_1-1} \right) \left(\sum_{j=k+1}^n \gamma_j^{a_1} \right)^{b_1-1} \left(\sum_{j=k+1}^n \gamma_j^{a_2} \right)^{b_2} \dots \left(\sum_{j=k+1}^n \gamma_j^{a_m} \right)^{b_m} \\ & \leq \gamma_{k+1} \left(\sum_{j=k+1}^n \gamma_j \right)^{a_1-1+a_1(b_1-1)+a_2 b_2+\dots+a_m b_m} = \gamma_{k+1} \left(\sum_{j=k+1}^n \gamma_j \right)^{r-1}. \end{aligned}$$

The desired conclusion follows.

LEMMA 3. Let $\psi(t) = (\lambda_{s+1} - t) \dots (\lambda_n - t)$, $\Phi(t) = (\lambda_s - t)^a \psi(t)$ where a is a positive integer, $0 \leq s < n$ and $\lambda_s < \lambda_{s+1}$. Then $\Phi^{(r)}(\lambda_s) = 0$ when $0 \leq r < a$, and when $r \geq a$,

$$|\Phi^{(r)}(\lambda_s)| \leq M \psi(\lambda_s) \left(\sum_{j=s+1}^n \frac{1}{\lambda_j - \lambda_s} \right)^{r-a}$$

where M is a number independent of s and n .

Proof. The first part is evident. For the second part we observe that, when $r \geq a$,

$$|\Phi^{(r)}(\lambda_s)| = r(r-1) \dots (r-a+1) \psi^{(r-a)}(\lambda_s),$$

and, as in the proof of Lemma 2, that $\psi^{(r-a)}(\lambda_s)/\psi(\lambda_s)$ can be expressed as a linear combination with constant coefficients of terms each with absolute value no greater than

$$\left(\sum_{j=s+1}^n \frac{1}{\lambda_j - \lambda_s} \right)^{r-a}.$$

The desired conclusion follows.

LEMMA 4. If $M_1 < \infty$, $\lambda_s < \lambda_{s+1}$ and $r = 0, 1, \dots, m_s - 1$, then

$$\mu_s^{(r)} = \lim_{n \rightarrow \infty} \sum_{k=s}^n \lambda_{nk} \left(1 - \frac{\lambda_s}{\lambda_{k+1}} \right) \dots \left(1 - \frac{\lambda_s}{\lambda_n} \right) \left(\sum_{j=k+1}^n \frac{1}{\lambda_j - \lambda_s} \right)^r.$$

Proof. For $r = 0$ the above sum is equal to $\mu_s^{(0)}$ for every $n \geq s$ by

(3). Suppose therefore that $1 \leq r \leq m_s - 1$. Then, by Lemmas 2 and 3 we have, for $n \geq s$, that

$$\begin{aligned} (11) \quad & \left| (-1)^r P_n^{(r)}(\lambda_s) - \sum_{k=s}^n \lambda_{nk} \left(1 - \frac{\lambda_s}{\lambda_{k+1}} \right) \dots \left(1 - \frac{\lambda_s}{\lambda_n} \right) \right. \\ & \left. \times \left(\sum_{j=k+1}^n \frac{1}{\lambda_j - \lambda_s} \right)^r \right| \\ & \leq M \sum_{k=s}^n |\lambda_{nk}| \frac{w_{nk}}{\lambda_{k+1} - \lambda_s} + M w_{ns} \sum_{k=s-m_s+1}^{s-1} |\lambda_{nk}| \end{aligned}$$

where M is a positive number independent of s and n , and

$$w_{nk} = \left(1 - \frac{\lambda_s}{\lambda_{k+1}} \right) \dots \left(1 - \frac{\lambda_s}{\lambda_n} \right) \left(\sum_{j=k+1}^n \frac{1}{\lambda_j - \lambda_s} \right)^{r-1}.$$

Since $\sum_{k=0}^n |\lambda_{nk}| \leq M_1$ for $n \geq 0$, and, by Lemma 1(i) and (ii), w_{nk} is uniformly bounded and $\lim_{n \rightarrow \infty} w_{nk} = 0$ for $k \geq s$, the right-hand side of (11) tends to 0 as $n \rightarrow \infty$. In view of (3), this establishes the desired conclusion.

LEMMA 5. If $M_1 < \infty$ and $r = 0, 1, \dots, m_s - 1$, then

$$(-1)^r \mu_s^{(r)} = \lim_{n \rightarrow \infty} \sum_{k=0}^n \lambda_{nk} \left(\frac{D_k}{D_n} \right)^{\lambda_s} \log^r \frac{D_k}{D_n}.$$

Proof. Suppose, without loss in generality, that $\lambda_s < \lambda_{s+1}$, and let

$$\delta_{nk} = \left(1 - \frac{\lambda_s}{\lambda_{k+1}} \right) \dots \left(1 - \frac{\lambda_s}{\lambda_n} \right) \left(\sum_{j=k+1}^n \frac{1}{\lambda_j - \lambda_s} \right)^r.$$

Then, by Lemma 1(ii) and (iii),

$$\lim_{n \rightarrow \infty} \sum_{k=0}^n \lambda_{nk} \left\{ \delta_{nk} - \left(\frac{D_k}{D_n} \right)^{\lambda_s} \log^r \frac{D_k}{D_n} \right\} = 0$$

since $\sum_{k=0}^n |\lambda_{nk}| \leq M_1$ for $n \geq 0$ and $D_n \rightarrow \infty$; and, by Lemma 1(ii) and Lemma 4,

$$\lim_{n \rightarrow \infty} \sum_{k=0}^n \lambda_{nk} \delta_{nk} = \mu_s^{(r)}.$$

The desired conclusion follows.

LEMMA 6. If a function $x \in BV$ is such that

$$\int_0^1 t^{\lambda_s} \log^r t \, dx(t) = 0 \quad \text{for } r = 0, 1, \dots, m_s - 1, \quad s = 0, 1, \dots,$$

then $x(t) = x(0+)$ for $0 < t \leq 1$. If, in addition, $\lambda_0 = 0$, then $x(0+) = 0$.

Proof. When $\lambda_0 = 0$ it follows from a known result (see [11, Theorem 8.2]) that

$$\int_0^1 t^n dx(t) = 0 \quad \text{for } n = 0, 1, \dots$$

The proof can now be completed in the same way as in the proof of Lemma 3 in [2].

3. Proofs of the main results.

Proofs of the necessity parts of Theorem 1(i), (ii) and (iii).

Part (i). Suppose the function $\alpha \in BV$ satisfies (1). For $0 \leq k \leq n$, we have that

$$\lambda_{nk} = \int_0^1 \lambda_{nk}(t) d\alpha(t),$$

and thus, by (6),

$$\sum_{k=0}^n |\lambda_{nk}| \leq \int_0^1 |d\alpha(t)| \sum_{k=0}^n \lambda_{nk}(t) \leq \int_0^1 |d\alpha(t)|.$$

Hence

$$(12) \quad M_1 \leq \int_0^1 |d\alpha(t)|.$$

Part (ii). Suppose the function $\beta \in L_\infty$ satisfies (8). For $0 \leq k \leq n$, we have that

$$(13) \quad \lambda_{nk} = \int_0^1 \lambda_{nk}(t) \beta(t) dt$$

and thus, by (6) and (7),

$$|\lambda_{nk}| \leq \|\beta\|_\infty \frac{d_k}{D_n}.$$

Hence

$$(14) \quad M_\infty \leq \|\beta\|_\infty.$$

Part (iii). Suppose the function $\beta \in L_\theta$ satisfies (8). It follows from (13) and (7), by Jensen's inequality (see [15, pp. 23-24]) that

$$\theta\left(\frac{D_n}{d_k} \lambda_{nk}\right) \leq \frac{D_n}{d_k} \int_0^1 \lambda_{nk}(t) \theta(\beta(t)) dt \quad \text{for } 0 \leq k \leq n.$$

Hence, by (6),

$$\sum_{k=0}^n \frac{d_k}{D_n} \theta\left(\frac{D_n}{d_k} \lambda_{nk}\right) \leq \int_0^1 \theta(\beta(t)) dt$$

and so

$$(15) \quad M_\theta \leq \int_0^1 \theta(\beta(t)) dt.$$

Proofs of the sufficiency parts of Theorem 1(i), (ii) and (iii).

We first observe that

$$\sum_{k=0}^n |\lambda_{nk}| \leq M_\infty \sum_{k=0}^n \frac{d_k}{D_n} \leq M_\infty,$$

and, by Young's inequality (see [8, p. 12]), that

$$\frac{D_n}{d_k} |\lambda_{nk}| \leq N(1) + \theta\left(\frac{D_n}{d_k} \lambda_{nk}\right)$$

where N is the convex function complementary to θ (see [8, p. 11]). Hence

$$\sum_{k=0}^n |\lambda_{nk}| \leq N(1) \sum_{k=0}^n \frac{d_k}{D_n} + \sum_{k=0}^n \frac{d_k}{D_n} \theta\left(\frac{D_n}{d_k} \lambda_{nk}\right) \leq N(1) + M_\theta.$$

It follows that $M_1 \leq M_\infty$, $M_1 \leq N(1) + M_\theta$ and so $M_1 < \infty$ under each of the three hypotheses of the sufficiency parts of Theorem 1(i), (ii) and (iii). Suppose therefore that $M_1 < \infty$.

For $n = 0, 1, \dots$, define the function α_n on $[0, 1]$ by setting

$$\alpha_n(t) = \begin{cases} 0 & \text{for } 0 \leq t < 1/D_n, \\ \sum_{D_k \leq t D_n} \lambda_{nk} & \text{for } 1/D_n \leq t \leq 1, \end{cases}$$

so that

$$\int_0^1 |d\alpha_n(t)| = \sum_{k=0}^n |\lambda_{nk}| \leq M_1.$$

Consequently, by Helly's theorem (see [14, p. 29]), there is an increasing sequence of positive integers $\{n_i\}$ and a function α of bounded variation on $[0, 1]$ such that

$$(16) \quad \lim_{i \rightarrow \infty} \alpha_{n_i}(t) = \alpha(t) \quad \text{for } 0 \leq t \leq 1$$

and

$$(17) \quad \int_0^1 |d\alpha(t)| \leq M_1.$$

Part (i). By Lemma 5, we have that

$$(-1)^r \mu_s^{(r)} = \lim_{n \rightarrow \infty} \sum_{k=0}^n \lambda_{nk} \left(\frac{D_k}{D_n}\right)^{\lambda_s} \log^r \frac{D_k}{D_n} = \lim_{n \rightarrow \infty} \int_0^1 t^{\lambda_s} \log^r t d\alpha_n(t)$$

for $r = 0, 1, \dots, m_s - 1$, $s = 0, 1, \dots$. It follows, by the Helly-Bray theorem, (see [14, p. 31]) that α satisfies (2) and hence (1).

Part (ii). Suppose $M_\infty < \infty$. Let $0 \leq x < y \leq 1$. Then for n sufficiently large there are integers a, b (depending on n) such that $-1 \leq a < b \leq n$ and

$$\frac{D_a}{D_n} \leq x < \frac{D_{a+1}}{D_n} \leq \frac{D_b}{D_n} \leq y < \frac{D_{b+1}}{D_n} \quad (D_{-1} = 0),$$

since

$$\max_{0 \leq k \leq n} \frac{d_k}{D_n} = \max_{0 \leq k \leq n} \frac{D_k}{D_n} \frac{1}{1 + \lambda_k} \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

Now

$$\frac{|\alpha_n(y) - \alpha_n(x)|}{\sum_{k=a+1}^b \frac{d_k}{D_n}} = \frac{\left| \sum_{k=a+1}^b \lambda_{nk} \right|}{\sum_{k=a+1}^b \frac{d_k}{D_n}} \leq M_\infty,$$

and

$$\lim_{n \rightarrow \infty} \sum_{k=a+1}^b \frac{d_k}{D_n} = y - x.$$

In view of (16), it follows that

$$\frac{|\alpha(y) - \alpha(x)|}{y - x} \leq M_\infty.$$

Hence

$$\alpha(t) = c + \int_0^t \beta(u) du \quad \text{for } 0 \leq t \leq 1$$

where $\beta \in L_\infty$ and $\|\beta\|_\infty \leq M_\infty$. Further, β satisfies (8) since α satisfies (1).

Part (iii). Suppose $M_\theta < \infty$. Let $0 = x_0 < x_1 < \dots < x_m = 1$. Then, for n sufficiently large, there exist integers a_0, a_1, \dots, a_m (depending on n) such that $-1 = a_0 < a_1 < \dots < a_m = n$ and

$$\frac{D_{a_j}}{D_n} \leq x_j < \frac{D_{1+a_j}}{D_n} \quad \text{for } j = 1, 2, \dots, m-1,$$

so that

$$\alpha_n(x_{j+1}) - \alpha_n(x_j) = \sum_{k=1+a_j}^{a_{j+1}} \lambda_{nk} \quad \text{for } j = 0, 1, \dots, m-1.$$

Let

$$\sigma_{jn} = \left(\sum_{k=1+a_j}^{a_{j+1}} \frac{d_k}{D_n} \right) \theta \left(\frac{\alpha_n(x_{j+1}) - \alpha_n(x_j)}{\sum_{k=1+a_j}^{a_{j+1}} \frac{d_k}{D_n}} \right).$$

Then, by Jensen's inequality (see [15, pp. 23-24]),

$$\sigma_{jn} \leq \sum_{k=1+a_j}^{a_{j+1}} \frac{d_k}{D_n} \theta \left(\frac{D_n}{d_k} \lambda_{nk} \right) \quad \text{for } j = 0, 1, \dots, m-1,$$

and so

$$\sum_{j=0}^{m-1} \sigma_{jn} \leq M_\theta.$$

Also

$$\lim_{n \rightarrow \infty} \sum_{k=1+a_j}^{a_{j+1}} \frac{d_k}{D_n} = x_{j+1} - x_j \quad \text{for } j = 0, 1, \dots, m-1.$$

In view of (16), it follows that

$$\lim_{n \rightarrow \infty} \sum_{j=0}^{m-1} \sigma_{jn} = \sum_{j=0}^{m-1} (x_{j+1} - x_j) \theta \left(\frac{\alpha(x_{j+1}) - \alpha(x_j)}{x_{j+1} - x_j} \right) \leq M_\theta,$$

and, by a theorem of Medvedev [12], this implies that

$$\alpha(t) = c + \int_0^t \beta(u) du \quad \text{for } 0 \leq t \leq 1$$

where $\beta \in L_\theta$ and $\int_0^1 \theta(\beta(t)) dt \leq M_\theta$. Further, β satisfies (8) since α satisfies (1).

Proofs of Theorem 1(iv), (v) and (vi).

Part (iv). Suppose that $l_0 = 0$. By Lemma 6 the function $\alpha \in BV$ satisfying (1) is unique. By (12), (17) and the proof of the sufficiency part of Theorem 1(i), we have that

$$M_1 \leq \int_0^1 |d\alpha(t)| \leq M_1.$$

Suppose that $l_0 > 0$, and let $\gamma(0) = 0$, $\gamma(t) = \alpha(t) - \alpha(0+)$ for $0 < t \leq 1$. Then $\gamma \in BV$ and satisfies (1). Hence, by (12),

$$M_1 \leq \int_0^1 |d\gamma(t)|.$$

Further, by (17) and the proof of the sufficiency part of Theorem 1(i), there is a function $\bar{\alpha} \in BV$ satisfying (1) and

$$\int_0^1 |d\bar{\alpha}(t)| \leq M_1.$$

By Lemma 6, $\gamma(t) = \bar{\alpha}(t) - \bar{\alpha}(0+)$ for $0 < t \leq 1$. Since $\gamma(0+) = \gamma(0)$, we have that

$$M_1 \leq \int_0^1 |d\gamma(t)| \leq \int_0^1 |d\bar{\alpha}(t)| \leq M_1.$$

Hence

$$M_1 = \int_0^1 |d\alpha(t)| - |\alpha(0+)|.$$

Part (v). By Lemma 6, the function $\beta \in L_\infty$ satisfying (8) is essentially unique. By (14) and the proof of the sufficiency part of Theorem 1(ii), we have that $M_\infty \leq \|\beta\|_\infty \leq M_\infty$.

Part (vi). This part can be established by the proof of Part (v) with certain obvious modifications.

Proof of Theorem 2. Let $0 \leq k \leq n$. Then

$$\begin{aligned} & \left(1 - \frac{\lambda_k}{\lambda_{n+1}}\right) \lambda_{n+1,k} + \frac{\lambda_{k+1}}{\lambda_{n+1}} \lambda_{n+1,k+1} \\ &= -\lambda_{k+1} \dots \lambda_{n+1} \left(1 - \frac{\lambda_k}{\lambda_{n+1}}\right) \frac{1}{2\pi i} \int_{c_{k,n+1}} \frac{P_{n+1}(z) dz}{(\lambda_k - z) \dots (\lambda_{n+1} - z)} \\ & \quad - \lambda_{k+2} \dots \lambda_{n+1} \frac{\lambda_{k+1}}{\lambda_{n+1}} \frac{1}{2\pi i} \int_{c_{k,n+1}} \frac{P_{n+1}(z) dz}{(\lambda_{k+1} - z) \dots (\lambda_{n+1} - z)} \\ &= -\lambda_{k+1} \dots \lambda_n \frac{1}{2\pi i} \int_{c_{k,n+1}} \frac{P_{n+1}(z) dz}{(\lambda_k - z) \dots (\lambda_n - z)} = \lambda_{nk}; \end{aligned}$$

and hence

$$(18) \quad \lambda_{nk} \frac{D_n}{d_k} = \left(1 - \frac{\lambda_k}{\lambda_{n+1}}\right) \lambda_{n+1,k} \frac{D_n}{d_k} + (1 + \lambda_k) \frac{\lambda_{n+1,k+1}}{\lambda_{n+1}} \frac{D_n}{d_{k+1}}.$$

It follows that

$$M_\infty(n) \leq M_\infty(n+1) \left(1 + \frac{1}{\lambda_{n+1}}\right) \frac{D_n}{D_{n+1}} = M_\infty(n+1).$$

Since

$$\left(1 - \frac{\lambda_k}{\lambda_{n+1}}\right) \frac{D_n}{D_{n+1}} + (1 + \lambda_k) \frac{D_n}{\lambda_{n+1} D_{n+1}} = 1,$$

applying Jensen's inequality to (18) yields

$$\begin{aligned} & \frac{d_k}{D_n} \theta \left(\frac{D_n}{d_k} \lambda_{nk} \right) \\ & \leq \frac{d_k}{D_n} \left\{ \left(1 - \frac{\lambda_k}{\lambda_{n+1}}\right) \frac{D_n}{D_{n+1}} \theta \left(\lambda_{n+1,k} \frac{D_{n+1}}{d_k} \right) \right. \\ & \quad \left. + (1 + \lambda_k) \frac{D_n}{\lambda_{n+1} D_{n+1}} \theta \left(\lambda_{n+1,k+1} \frac{D_{n+1}}{d_{k+1}} \right) \right\} \\ & = \left(1 - \frac{\lambda_k}{\lambda_{n+1}}\right) \frac{d_k}{D_{n+1}} \theta \left(\lambda_{n+1,k} \frac{D_{n+1}}{d_k} \right) + \frac{\lambda_{k+1} d_{k+1}}{\lambda_{n+1} D_{n+1}} \theta \left(\lambda_{n+1,k+1} \frac{D_{n+1}}{d_{k+1}} \right). \end{aligned}$$

Summing this inequality for $k = 0, 1, \dots, n$, we get that

$$M_\theta(n) \leq M_\theta(n+1) - \frac{\lambda_0 d_0}{\lambda_{n+1} D_{n+1}} \theta \left(\lambda_{n+1,0} \frac{D_{n+1}}{d_0} \right) \leq M_\theta(n+1).$$

Since the above argument is valid when θ is any even continuous convex function, we can take $\theta(u) = |u|$ to obtain, in addition, that

$$M_1(n) \leq M_1(n+1).$$

This completes the proof of Theorem 2.

Note. In all but Theorem 2 the condition that the sequences $\{l_n\}$ and $\{\lambda_n\}$ be monotonic is redundant and was imposed only to avoid non-essential and tedious complication in the proofs. Without the monotonicity condition, but with $\{l_n\}$ distinct, $\lambda_0 = l_0 \geq 0$, $k_0 = 1$ if $l_0 = 0$, $l_n > 0$ for $n = 1, 2, \dots$, identities and inequalities such as (5), (6) (using (10) and (11) on p. 46 of [11] and the proof of Lemma 1 in [10]) and (7) can readily be shown to hold, and Lemmas 5 and 6 and Theorem 1 remain valid. Removal of the monotonicity condition involves changes in statements and proofs of lemmas as indicated below.

Statements.

LEMMA 1. Replace $0 < \lambda < \lambda_{a+1}$ by $0 < \lambda < \min_{k>a} \lambda_k$.

LEMMA 2. Replace $0 < t < \lambda_{k+1}$ by $0 < t \neq \lambda_j$ for $j > k$, and

$$\frac{\psi(t)}{\lambda_{k+1} - t} \left(\sum_{j=k+1}^n \frac{1}{\lambda_j - t} \right)^{r-1} \text{ by } \max_{i>k} \frac{|\psi(t)|}{|\lambda_i - t|} \left(\sum_{j=k+1}^n \frac{1}{|\lambda_j - t|} \right)^{r-1}.$$

LEMMA 3. Replace $\lambda_s < \lambda_{s+1}$ by $\lambda_s \neq \lambda_j$ for $n > j > s$, and

$$\psi(\lambda_s) \left(\sum_{j=s+1}^n \frac{1}{\lambda_j - \lambda_s} \right)^{r-a} \text{ by } |\psi(\lambda_s)| \left(\sum_{j=s+1}^n \frac{1}{|\lambda_j - \lambda_s|} \right)^{r-a}.$$

LEMMA 4. Replace $\lambda_s < \lambda_{s+1}$ by $\lambda_s \neq \lambda_j$ for $j > s$.

Proofs.

Lemma 1. Replace $\lambda_{k+1}/(\lambda_{k+1} - \lambda)$ by $\max_{j>k} \lambda_j/(\lambda_j - \lambda)$, and $1/\lambda_{k+1}$ by $\max_{j>k} 1/\lambda_j$.

Lemma 2. In the inequalities replace γ_j by $|\gamma_j|$ and γ_{k+1} by $\max_{j>k} |\gamma_j|$.

Lemma 3. Replace $\lambda_j - \lambda_s$ by $|\lambda_j - \lambda_s|$.

Lemma 4. Replace $1/(\lambda_{k+1} - \lambda_s)$ by $\max_{j>k} 1/|\lambda_j - \lambda_s|$, and take

$$w_{nk} = \left| \left(1 - \frac{\lambda_s}{\lambda_{k+1}}\right) \dots \left(1 - \frac{\lambda_s}{\lambda_n}\right) \right| \left(\sum_{j=k+1}^n \frac{1}{|\lambda_j - \lambda_s|} \right)^{r-1}.$$

Lemma 5. Replace $\lambda_s < \lambda_{s+1}$ by $\lambda_s \neq \lambda_j$ for $j > s$.

REFERENCES

1. D. L. Berman, *Application of interpolatory polynomial operators to solve the moment problem*, *Urkain Math. Z.* 14 (1963), 184-190.
2. D. Borwein, *The Hausdorff moment problem*, *Can. Math. Bull.* 21 (1978), 257-265; *Corrections* 22 (1979), 128.
3. P. J. Davis, *Interpolation and approximation* (Blaisdell, New York, 1965).
4. K. Endl, *On systems of linear inequalities in infinitely many variables and generalized Hausdorff means*, *Math. Z.* 82 (1963), 1-7.
5. F. Hausdorff, *Summationsmethoden und Momentfolgen I*, *Math. Z.* 9 (1921), 74-109.
6. ——— *Summationsmethoden und Momentfolgen II*, *Math. Z.* 9 (1921), 280-299.
7. ——— *Momentprobleme für ein endliches Intervall*, *Math. Z.* 16 (1923), 220-248.
8. M. A. Krasnosel'skiĭ and Ya. B. Rutickii, *Convex functions and Orlicz spaces* (P. Noordhoff, Groningen, 1961).
9. D. Leviatan, *A generalized moment problem*, *Israel J. Math.* 5 (1967), 97-103.
10. ——— *Some moment problems in a finite interval*, *Can. J. Math.* 20 (1968), 960-966.
11. G. G. Lorentz, *Bernstein polynomials* (University of Toronto Press, Toronto, 1953).
12. U. T. Medvedev, *Generalization of a theorem of F. Riesz*, *Uspeshi, Mat. Nauk* 8 (1953), 115-118.
13. I. J. Schoenberg, *On finite rowed systems of linear inequalities in infinitely many variables*, *Trans. A.M.S.* 34 (1932), 594-619.
14. D. V. Widder, *The Laplace transform* (Princeton, 1946).
15. A. Zygmund, *Trigonometric series I*, 2nd edition (Cambridge University Press, Cambridge, 1959).

*The University of Western Ontario,
London, Ontario;
Tel-Aviv University,
Tel-Aviv, Israel*