

## NÖRLUND OPERATORS ON $l_p$

DAVID BORWEIN

ABSTRACT. The Nörlund matrix  $N_a$  is the triangular matrix  $\{a_{n-k}/A_n\}$ , where  $a_n \geq 0$  and  $A_n := a_0 + a_1 + \cdots + a_n > 0$ . It is proved that, subject to the existence of  $\alpha := \lim_{n \rightarrow \infty} na_n/A_n$ ,  $N_\alpha \in B(l_p)$  for  $1 < p < \infty$  if and only if  $\alpha < \infty$ . It is also proved

REPRINTED FROM / RÉIMPRIMÉ DE

---

Canadian Mathematical

# Bulletin

canadien de mathématiques

---



The Canadian Mathematical Society

La Société mathématique du Canada

## NÖRLUND OPERATORS ON $l_p$

DAVID BORWEIN

**ABSTRACT.** The Nörlund matrix  $N_a$  is the triangular matrix  $\{a_{n-k}/A_n\}$ , where  $a_n \geq 0$  and  $A_n := a_0 + a_1 + \dots + a_n > 0$ . It is proved that, subject to the existence of  $\alpha := \lim na_n/A_n$ ,  $N_a \in B(l_p)$  for  $1 < p < \infty$  if and only if  $\alpha < \infty$ . It is also proved that it is possible to have  $N_a \in B(l_p)$  for  $1 < p < \infty$  when  $\sup na_n/A_n = \infty$ .

**1. Introduction.** Let  $a := \{a_n\}$  be a sequence of non-negative numbers, and let  $A_n := a_0 + a_1 + \dots + a_n > 0$ . The Nörlund matrix  $N_a := \{a_{nk}\}$  is defined by

$$a_{nk} := \begin{cases} a_{n-k}/A_n & \text{for } 0 \leq k \leq n, \\ 0 & \text{for } k > n. \end{cases}$$

The  $N_a$ -transform  $y = \{y_n\}$  of the sequence  $x = \{x_n\}$  is given by

$$y_n := (N_a x)_n := \frac{1}{A_n} \sum_{k=0}^n a_{n-k} x_k.$$

Suppose throughout that

$$1 < p < \infty, \quad \frac{1}{p} + \frac{1}{q} = 1,$$

and define

$$\begin{aligned} \sigma_1(n) &:= \frac{1}{A_n} \sum_{k=0}^n a_{n-k} \left(\frac{n+1}{k+1}\right)^{1/p}, \\ \sigma_2(k) &:= \sum_{n=k}^{\infty} \frac{a_{n-k}}{A_n} \left(\frac{k+1}{n+1}\right)^{1/q}, \\ M_1 &:= \sup_{n \geq 0} \sigma_1(n), \quad M_2 := \sup_{k \geq 0} \sigma_2(k). \end{aligned}$$

Let

$$\|N_a\|_p := \sup_{\|x\|_p \leq 1} \|N_a x\|_p,$$

where

$$\|x\|_p := \left( \sum_{n=0}^{\infty} |x_n|^p \right)^{1/p},$$

so that  $N_a \in B(l_p)$ , the Banach algebra of bounded linear operators on  $l_p$ , exactly when  $\|N_a\|_p$  is finite (in which case it is the norm of  $N_a$ ).

The following theorem concerning sufficient conditions for  $N_a \in B(l_p)$  is due to Borwein and Cass [1, Theorem 2].

This research was supported in part by the Natural Sciences and Engineering Research Council of Canada.

Received by the editors June 3, 1991; revised November 27, 1991.

AMS subject classification: Primary: 47B37, 47A30; secondary: 40G05.

Key words and phrases: Nörlund, operators on  $l_p$ , norm estimates.

© Canadian Mathematical Society 1993.



THEOREM A. If

$$(1) \quad \frac{na_n}{A_n} = O(1),$$

then  $N_a \in B(l_p)$  and  $\|N_a\|_p \leq M_1^{1/q} M_2^{1/p} < \infty$ .

Cass and Kratz [3] showed that (1) is in fact necessary and sufficient for  $N_a \in B(l_p)$  when

$$(2) \quad a_n := f(n),$$

where  $f(x)$  is a logarithmico-exponential function for all sufficiently large positive values of  $x$ . They showed that  $na_n/A_n$  tends to a finite or infinite limit when (2) is satisfied, and proved:

THEOREM B. Suppose that  $a_n$  is given by (2), and that  $na_n/A_n \rightarrow \alpha$ .

(i) Then  $N_a \in B(l_p)$  if and only if  $\alpha < \infty$ .

(ii) If  $\alpha < \infty$ , then

$$\lim_{n \rightarrow \infty} \sigma_1(n) = \frac{\Gamma(\alpha + 1)\Gamma(1/q)}{\Gamma(\alpha + 1/q)} \leq \|N_a\|_p \leq M_1^{1/q} M_2^{1/p} < \infty.$$

Condition (2) is redundant when  $\alpha = 0$ .

For  $\alpha > -1$ , the Cesàro matrix  $C_\alpha$  is the Nörlund matrix  $N_a$  with

$$a_n := \binom{n + \alpha - 1}{n}.$$

It follows from an inequality of Hardy's [4] that, for  $\alpha \geq 0$ ,  $C_\alpha \in B(l_p)$  and

$$\|C_\alpha\|_p = \frac{\Gamma(\alpha + 1)\Gamma(1/q)}{\Gamma(\alpha + 1/q)}.$$

This is thus the attained lower bound of the norms of all Nörlund matrices satisfying the conditions of Theorem B(ii).

The primary object of this paper is to show that the requirement that  $a_n$  be generated by a logarithmico-exponential function is redundant in Theorem B(i), and can be replaced by a far less restrictive monotonicity condition in Theorem B(ii). To this end we shall prove:

THEOREM 1. Suppose that  $na_n/A_n \rightarrow \alpha$ .

(i) Then  $N_a \in B(l_p)$  if and only if  $\alpha < \infty$ .

(ii) If  $\alpha < \infty$ , then

$$\|N_a\|_p \leq M_1^{1/q} M_2^{1/p} < \infty,$$

and if, in addition,  $\{n^c a_n\}$  is eventually monotonic for every constant  $c \neq 1 - \alpha$ , then

$$\lim_{n \rightarrow \infty} \sigma_1(n) = \frac{\Gamma(\alpha + 1)\Gamma(1/q)}{\Gamma(\alpha + 1/q)} \leq \|N_a\|_p.$$

Further, the monotonicity condition is redundant when  $\alpha = 0$ .

In § 4 we construct a sequence  $\{a_n\}$  such that  $\{na_n/A_n\}$  is unbounded but  $N_a \in B(l_p)$  for all finite  $p > 1$ , thus showing that (1) is not a necessary condition for  $N_a \in B(l_p)$ .

ACKNOWLEDGEMENT. I am indebted to Xiaopeng Gao, a graduate student at the University of Western Ontario, for the removal of a redundant monotonicity condition from Theorem 1 and Lemma 1 (below). I had proved these results subject to  $\{a_n/A_n\}$  being eventually monotonic. Gao's incisive analysis of my original proofs showed how they could be adjusted to establish the present more satisfactory forms of the results.

## 2. Preliminary results.

LEMMA 1. Suppose that  $\{a_n/A_n\}$  is positive and null. Let

$$h := h(n) := \max_{k \leq n} \left[ \frac{A_k}{a_k} \right].$$

Then

$$A_n \geq \sum_{n-h \leq k \leq n} a_k \geq \left(1 - \frac{1}{e}\right) A_n.$$

Further, if  $na_n/A_n \rightarrow \infty$ , then  $h(n) = o(n)$ .

PROOF. Suppose without loss that  $n - h \geq 1$ . Then

$$\begin{aligned} \log \frac{A_n}{A_{n-h-1}} &= \int_{A_{n-h-1}}^{A_n} \frac{dx}{x} = \sum_{k=n-h}^n \int_{A_{k-1}}^{A_k} \frac{dx}{x} \geq \sum_{k=n-h}^n \frac{a_k}{A_k} \\ &\geq (h+1) \min_{k \leq n} \frac{a_k}{A_k} \geq 1, \end{aligned}$$

and so

$$\frac{A_{n-h-1}}{A_n} \leq \frac{1}{e}.$$

Thus

$$\frac{1}{A_n} \sum_{k=n-h}^n a_k = 1 - \frac{A_{n-h-1}}{A_n} \geq 1 - \frac{1}{e}.$$

Now suppose that  $na_n/A_n \rightarrow \infty$ . Then, for  $m \geq 1$ ,

$$\begin{aligned} 0 &\leq \limsup_{n \rightarrow \infty} \frac{h(n)}{n} \leq \lim_{n \rightarrow \infty} \frac{1}{n} \max_{k \leq m} \frac{A_k}{a_k} + \limsup_{n \rightarrow \infty} \max_{m \leq k \leq n} \frac{k}{n} \frac{A_k}{ka_k} \\ &\leq \max_{k \geq m} \frac{A_k}{ka_k} \rightarrow 0 \text{ as } m \rightarrow \infty. \end{aligned}$$

It follows that  $h(n)/n \rightarrow 0$ .

LEMMA 2. Suppose that  $na_n/A_n \rightarrow \alpha$  where  $0 < \alpha < \infty$ . Then

$$\lim_{n \rightarrow \infty} n^c a_n = \begin{cases} \infty & \text{if } c > 1 - \alpha, \\ 0 & \text{if } c < 1 - \alpha. \end{cases}$$

PROOF. Let

$$\alpha_n := \frac{na_n}{A_n}.$$

Then, as  $n \rightarrow \infty$ ,

$$\log A_n - \log A_{n-1} = -\log\left(1 - \frac{\alpha_n}{n}\right) \sim \frac{\alpha_n}{n},$$

and hence

$$\log A_n = \log A_0 - \sum_{k=1}^{n-1} \log\left(1 - \frac{\alpha_k}{k}\right) = (\alpha + o(1)) \log n.$$

Consequently  $A_n \sim n^{\alpha+o(1)}$ , and so  $a_n \sim \alpha n^{\alpha-1+o(1)}$ . The desired conclusion follows. ■

LEMMA 3. Suppose that  $0 < \alpha < \infty$ ,  $\delta < 1$ , and that  $n^c a_n$  is eventually positive and increasing when the constant  $c > 1 - \alpha$ , and eventually decreasing when  $c < 1 - \alpha$ . Then, as  $n \rightarrow \infty$ ,

$$I(n) := \frac{1}{n} \sum_{k=0}^n \frac{a_k}{a_n} \left(1 - \frac{k}{n+1}\right)^{-\delta} \rightarrow \frac{\Gamma(\alpha)\Gamma(1-\delta)}{\Gamma(\alpha+1-\delta)}.$$

PROOF. Let  $1 > c_1 > 1 - \alpha > c_2$ . Then there is a positive integer  $N$  such that

$$\left(\frac{k}{n}\right)^{-c_2} \leq \frac{a_k}{a_n} \leq \left(\frac{k}{n}\right)^{-c_1} \text{ for } n \geq k \geq N.$$

Since

$$\lim_{n \rightarrow \infty} na_n = \infty,$$

we obtain

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=0}^{N-1} \frac{a_k}{a_n} \left(1 - \frac{k}{n+1}\right)^{-\delta} = 0,$$

and hence

$$\begin{aligned} \limsup_{n \rightarrow \infty} I(n) &\leq \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=N}^n \left(\frac{k}{n}\right)^{-c_1} \left(1 - \frac{k}{n+1}\right)^{-\delta} \\ &= \int_0^1 x^{-c_1} (1-x)^{-\delta} dx = \frac{\Gamma(1-c_1)\Gamma(1-\delta)}{\Gamma(2-c_1-\delta)} \end{aligned}$$

and

$$\begin{aligned} \liminf_{n \rightarrow \infty} I(n) &\geq \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=N}^n \left(\frac{k}{n}\right)^{-c_2} \left(1 - \frac{k}{n+1}\right)^{-\delta} \\ &= \int_0^1 x^{-c_2} (1-x)^{-\delta} dx = \frac{\Gamma(1-c_2)\Gamma(1-\delta)}{\Gamma(2-c_2-\delta)}. \end{aligned}$$

Letting  $c_1 \rightarrow 1 - \alpha$  from the right and  $c_2 \rightarrow 1 - \alpha$  from the left, we get the desired conclusion. ■

The following lemma is a special case of a known result [2, Theorems 2].

LEMMA 4. If  $\{b_n\}$  is a sequence of positive numbers, and if

$$\mu_1 := \sup_{n \geq 0} \sum_{k=0}^n \frac{a_{n-k}}{A_n} \left(\frac{b_k}{b_n}\right)^{1/p} < \infty \text{ and } \mu_2 := \sup_{k \geq 0} \sum_{n=k}^{\infty} \frac{a_{n-k}}{A_n} \left(\frac{b_n}{b_k}\right)^{1/q} < \infty,$$

then  $N_a \in B(l_p)$  and  $\|N_a\|_p \leq \mu_1^{1/q} \mu_2^{1/p}$ .

LEMMA 5. If  $\{b_n\}$  is a bounded sequence of positive numbers such that  $\sum b_n = \infty$ , and if, as  $n \rightarrow \infty$ ,

$$\sum_{k=0}^n \frac{a_{n-k}}{A_n} \left(\frac{b_k}{b_n}\right)^{1/p} \rightarrow \sigma \text{ (finite or infinite),}$$

then  $\|N_a\|_p \geq \sigma$ .

PROOF. Observe that if

$$D_n := \prod_{k=0}^n \left(1 - \frac{b_k}{b}\right)^{-1} \text{ where } b > \sup_{k \geq 0} b_k,$$

and  $d_n := D_n - D_{n-1}$  for  $n \geq 1$ , then  $b_n = b d_n / D_n$  and  $D_n \rightarrow \infty$ . The desired conclusion is now a consequence of a known result [2, Theorem 4]. ■

### 3. Proof of Theorem 1.

PART (i). In view of Theorem A, it suffices to show that  $N_a \notin B(l_p)$  when  $\alpha = \infty$ . Suppose therefore that  $\alpha = \infty$ .

If  $\limsup a_n/A_n > 0$ , then  $\sum |a_n/A_n|^p = \infty$ , but this implies that  $N_a e^0 \notin l_p$ , where  $e^0 = (1, 0, 0, \dots)$ , so that  $N_a \notin B(l_p)$ .

Suppose that  $\lim a_n/A_n = 0$ . Since  $\alpha = \infty$ , we have, by Lemma 1 with  $\delta := 1/p$ , that, as  $n \rightarrow \infty$ ,

$$\begin{aligned} \sigma_1(n) &= \frac{(n+1)^\delta}{A_n} \sum_{k=0}^n a_k (n+1-k)^{-\delta} \geq \frac{(n+1)^\delta}{A_n} \sum_{n-h \leq k \leq n} a_k (n+1-k)^{-\delta} \\ &\geq \left(\frac{n+1}{h+1}\right)^\delta \frac{1}{A_n} \sum_{n-h \leq k \leq n} a_k \geq \left(1 - \frac{1}{e}\right) \left(\frac{n+1}{h+1}\right)^\delta \rightarrow \infty. \end{aligned}$$

It follows, by Lemma 5, that  $N_a \notin B(l_p)$ .

PART (ii). The case  $\alpha = 0$  is part of Theorem B, so suppose that  $0 < \alpha < \infty$ . Since  $na_n = O(A_n)$ , it follows from Theorem A that  $N_a \in B(l_p)$  and  $\|N_a\|_p \leq M_1^{1/q} M_2^{1/p} < \infty$ .

The monotonicity condition together with Lemma 2 ensures that  $a_n$  satisfies the monotonicity conditions of Lemma 3. Hence, by Lemma 3 with  $\delta := 1/p$ ,

$$\sigma_1(n) = \frac{na_n}{A_n} I(n) \rightarrow \frac{\Gamma(\alpha+1)\Gamma(1/q)}{\Gamma(\alpha+1/q)}$$

and so, by Lemma 5,

$$\|N_a\|_p \geq \frac{\Gamma(\alpha+1)\Gamma(1/q)}{\Gamma(\alpha+1/q)}. \quad \blacksquare$$



4. **Construction.** We construct a sequence  $\{a_n\}$  such that  $\{na_n/A_n\}$  is unbounded but  $N_a \in B(l_p)$  for all finite  $p > 1$ . Let

$$a_n := \begin{cases} 1 & \text{when } n = m^2, m = 0, 1, \dots, \\ 0 & \text{otherwise.} \end{cases}$$

Then, for  $n = m^2$ ,

$$\frac{na_n}{A_n} = \frac{m^2}{m+1} \rightarrow \infty \text{ as } m \rightarrow \infty.$$

Thus  $\{na_n/A_n\}$  is unbounded.

Now define

$$S_1(n) := \sum_{k=0}^n \frac{a_{n-k}}{A_n} \left(\frac{b_k}{b_n}\right)^{1/p}, \quad S_2(k) := \sum_{n=k}^{\infty} \frac{a_{n-k}}{A_n} \left(\frac{b_n}{b_k}\right)^{1/q},$$

where

$$b_n := \frac{1}{\sqrt{n+1}}.$$

In order to prove that  $N_a \in B(l_p)$  it suffices, by Lemma 4, to demonstrate that  $S_1(n) = O(1)$  and  $S_2(k) = O(1)$ .

For  $1 \leq m^2 \leq n < (m+1)^2$ ,  $\delta := 1/(2p)$ , we have

$$\begin{aligned} S_1(n) &= \frac{(n+1)^\delta}{A_n} \sum_{k=0}^n a_k (n+1-k)^{-\delta} = \frac{(n+1)^\delta}{m+1} \sum_{k=0}^m (n+1-k^2)^{-\delta} \\ &\leq \frac{(n+1)^\delta (n+1-m^2)^{-\delta}}{m+1} + \frac{1}{m+1} + \frac{(n+1)^\delta}{m+1} \sum_{k=1}^{m-1} (m^2-k^2)^{-\delta} \\ &\leq (m+1)^{2\delta-1} + (m+1)^{-1} + (m+1)^{2\delta-1} m^{-\delta} \sum_{k=1}^{m-1} (m-k)^{-\delta} \\ &= (m+1)^{2\delta-1} + (m+1)^{-1} + (m+1)^{2\delta-1} m^{-\delta} \sum_{k=1}^{m-1} k^{-\delta} \\ &= O(m^{2\delta-1} + m^{-1} + m^{2\delta-1} \cdot m^{1-2\delta}) = O(1). \end{aligned}$$

Further, for  $1 \leq m^2 \leq k+1 < (m+1)^2$ ,  $\mu := 1/(2q)$ , we have

$$\begin{aligned} S_2(k) &= (k+1)^\mu \sum_{n=k}^{\infty} \frac{a_{n-k}}{A_n (n+1)^\mu} \\ &\leq (k+1)^\mu \sum_{n=k}^{2k} \frac{a_{n-k}}{A_n (n+1)^\mu} + (k+1)^\mu \sum_{n=k+1}^{\infty} \frac{a_n}{A_n (n+1)^\mu} \\ &\leq 1 + (m+1)^{2\mu} \sum_{r=m}^{\infty} \frac{1}{r^{1+2\mu}} = 1 + O(m^{2\mu} \cdot m^{-2\mu}) = O(1). \end{aligned}$$

Hence  $N_a \in B(l_p)$ .

## REFERENCES

1. D. Borwein and F. P. Cass, *Nörlund matrices as bounded operators on  $l_p$* , *Archiv der Mathematik* **42**(1984), 464-469.
2. D. Borwein and A. Jakimovski, *Matrix operators on  $l_p$* , *Rocky Mountain J. Math.* **9**(1979), 463-477.
3. F. P. Cass and W. Kratz, *Nörlund and weighted mean matrices as bounded operators on  $l_p$* , *Rocky Mountain J. Math.* **20**(1990), 59-74.
4. G. H. Hardy, *An inequality for Hausdorff means*, *J. London Math. Soc.* **18**(1943), 46-50.

Department of Mathematics  
University of Western Ontario  
London, Ontario  
N6A 5B7